

HERPES SIMPLEX VIRUS -HSV II-IGM

Profile Details | 1 Parameters

Fasting: Fasting is not mandatory | Report Time: Same Day



Specimen Required: Blood

Price- ₹780 **₹425**

Parameters Included

About This Test

A Herpes simplex virus II (HSV-2) IgM test measures the presence of IgM antibodies against HSV-2 in the blood. Here's what this entails: - **HSV-2:** This virus primarily causes genital herpes, characterized by sores and blisters in the genital and anal areas. It can also cause infections in other parts of the body. - **IgM Antibodies:** Immunoglobulin M (IgM) is one of the first types of antibodies produced by the immune system in response to an infection. The presence of IgM antibodies generally indicates a recent or active infection. - **Test Interpretation:** - **Positive HSV-2 IgM:** Suggests a recent or current HSV-2 infection. However, IgM tests can sometimes produce false positives and do not distinguish well between primary (initial) and recurrent infections. - **Negative HSV-2 IgM:** Suggests that there is no recent or current HSV-2 infection, although it does not rule out past infection. - **Limitations:** The HSV-2 IgM test can have limitations, including cross-reactivity with HSV-1, leading to potential false positives. Additionally, because IgM levels can decline quickly after the initial infection, they may not be present during later stages or recurrent infections. Overall, the HSV-2 IgM test is used to detect recent or active HSV-2 infections and is often part of a broader diagnostic approach, which may include IgG antibody tests and clinical evaluation.



98% Reports
within 24 hours



95% Samples
Delhi Accredited Labs



Home Collection
Door-to-door pickup



700+ Centres
Across India



750+ Tests
Available Profiles

**Fasting 10-12 hrs essential | *Limited Period Offer | *Reports within Same Day*

For appointments and queries: **82733 39996** | www.thexpertlab.com